

IMPORTANT RECOMMENDATION ON ENDOSCOPIC TRAINING AND QUALITY ASSURANCE

As professional organizations we strongly recommend that each hospital teaching endoscopic surgery should make available an endoscopic dry lab for training and improving the proficiency of the endoscopic surgery skills of the physician.

Evidence is provided that dry lab training prior to training in the OR, reduces patients' morbidity and mortality in all endoscopic surgical disciplines, and as such the skill testing infrastructure should aim for multi-disciplinary usage.

ESGE EBCOG EAGS ENTOG ACOG **AAGL** European **+heAcademy** of Gvnaecological Surgery Rudi Campo Chiara Benedetto **Arnaud Wattiez** Maud van de Venne Hal Lawrence Ceana H. Nezhat We change my Corale Northal ILD.

April 8, 2014

- 1. Palter VN, Orzech N, Reznick RK, Grantcharov TP. Validation of a structured training and assessment curriculum for technical skill acquisition in minimally invasive surgery: a randomized controlled trial. Ann Surg. 2013 Feb;257(2):224-30.
- 2. Gala R, Orejuela F, Gerten K, Lockrow E, Kilpatrick C, Chohan L, Green C, Vaught J, Goldberg A, Schaffer J. Effect of validated skills simulation on operating room performance in obstetrics and gynecology residents: a randomized controlled trial. Obstet Gynecol. 2013 Mar;121(3):578-84.
- 3. Molinas CR, Campo R. Defining a structured training program for acquiring basic and advanced laparoscopic psychomotor skills in a simulator. Gynecol Surg. 2010 Nov;7(4):427-35.
- 4. Burden C, Oestergaard J, Larsen CR. Integration of laparoscopic virtual-reality simulation into gynaecology training. BJOG. 2011 Nov;118 Suppl 3:5-10.
- 5. Hur HC, Arden D, Dodge LE, Zheng B, Ricciotti HA. Fundamentals of laparoscopic surgery: a surgical skills assessment tool in gynecology. JSLS. 2011 Jan-Mar;15(1):21-6.
- Campo R, Molinas CR, De Wilde RL, Brolmann H, Brucker S, Mencaglia L, Odonovan P, Wallwiener D, Wattiez A. Are you good enough for your patients? The European certification model in laparoscopic surgery. Facts Views Vis Obgyn. 2012;4(2):95-101.

